

Purified Anti-Human CD56/NCAM Antibody[B-A19]

Catalog No. E-AB-F1305A

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone	B-A19
Conjugation	Unconjugated
Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended Dilution

FCM	2 $\mu\text{g/mL}$ (1×10^5 – 5×10^5 cells)
------------	---------------------------------------------------------------

Data



Human peripheral blood lymphocytes were stained with 0.2 μg Purified Anti-Human CD56 Antibody[B-A19] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control(Left), followed by FITC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage This product can be stored at 2–8°C for 12 months. Do not freeze.

Background

CD56 is a single transmembrane glycoprotein also known as NCAM (neural cell adhesion molecule), Leu-19, or NKH1. It is a member of the Ig superfamily. The 140 kD isoform is expressed on NK and NKT cells. CD56 is also expressed in the brain (cerebellum and cortex) and at neuromuscular junctions. Certain large granular lymphocyte (LGL) leukemias, small-cell lung carcinomas, neuronal derived tumors, myelomas, and myeloid leukemias also express CD56. CD56 plays a role in homophilic and heterophilic adhesion via binding to itself or heparan sulfate.

For Research Use Only